

AD-A161 270

REAL INVERSION FORMULAS FOR LAPLACE AND STIELTJES
TRANSFORMS(U) NORTH CAROLINA UNIV AT CHAPEL HILL CENTER
FOR STOCHASTIC PROC. J L TEUGELS JUL 85 TR-111

1/1

UNCLASSIFIED

AFOSR-TR-85-0949 F49620-82-C-0009


F/G 12/1

NL

END

FILMED

DTIC




MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A161 270

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



REAL INVERSION FORMULAS FOR LAPLACE AND STIELTJES TRANSFORMS

by

Jozef L. Teugels

Technical Report No. 111

July 1985

DTIC
ELECTE
NOV 21 1985
S B D

DTIC FILE COPY

Approved for public release,
distribution unlimited

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT UNLIMITED	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 85-03949	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Rept. No. 111			7a. NAME OF MONITORING ORGANIZATION	
6a. NAME OF PERFORMING ORGANIZATION Center for Stochastic Processes		6b. OFFICE SYMBOL (If applicable)		7b. ADDRESS (City, State and ZIP Code) AFOSR/NM Bldg 410 Bolling AFB DC 20332-6448
6c. ADDRESS (City, State and ZIP Code) Statistics Dept., 321 PH 039A, UNC Chapel Hill, NC 27514		8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		
8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620-82-C-0009		
8c. ADDRESS (City, State and ZIP Code) Bolling AFB Washington, DC 20332		10. SOURCE OF FUNDING NOS.		
11. TITLE (Include Security Classification) REAL INVERSION FORMULAS FOR LAPLACE AND STIELTJES TRANSFORMS		PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304	TASK NO. A5
12. PERSONAL AUTHOR(S) Jozef L. Teugels		13. TYPE OF REPORT technical		
13b. TIME COVERED FROM 9/84 TO 8/85		14. DATE OF REPORT (Yr., Mo., Day) July 1985		15. PAGE COUNT 16
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Inversion formulas, Laplace transform, Stieltjes transform		
FIELD	GROUP	SUB. GR.		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) We provide probabilistic proofs for a number of real inversion formulas for the Laplace and for the Stieltjes transform.				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Brian W. Woodruff		22b. TELEPHONE NUMBER (Include Area Code) 919 702-2907		22c. OFFICE SYMBOL NM

DD FORM 147 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE

85 11 15 050

REAL INVERSION FORMULAS FOR LAPLACE AND STIELTJES TRANSFORMS

by

Jozef L. Teugels

Katholieke Universiteit te Leuven

and

University of North Carolina at Chapel Hill

Abstract

We provide probabilistic proofs for a number of real inversion formulas for the Laplace and for the Stieltjes transform.

Keywords: Inversion formulas, Laplace transform, Stieltjes transform

This research supported in part by the Air Force Office of Scientific Research, Contract No. F49620 82 C 0009, Center for Stochastic Processes, University of North Carolina, Chapel Hill, NC.

1. Introduction

Assume μ to be a bounded measure on \mathbb{R}_+ . Let f be a kernel changing μ into the transformed function

$$(1) \quad g(t) := \int_0^{\infty} f(t,u) d\mu(u).$$

In the literature dealing with operational calculus the recovery of μ from g is mostly accomplished by complex inversion formulas. One knows that for a variety of integral transforms (i.e. Laplace, Stieltjes), real inversion formulas are available [6,10,12]. These formulas can also be obtained as applications of classical results from probability theory such as the weak law of large numbers and the central limit theorem.

To abbreviate the writing we put for $0 < y_1 < y_2 < \infty$

$$(2) \quad \mu\{y_1; y_2\} = \frac{1}{2}\mu\{y_1\} + \mu(y_1, y_2) + \frac{1}{2}\mu\{y_2\}$$

Accession No.

DTIS 9-185

DTIC 118

Unannounced

Justification

BY

Distribution

Availability

Dist Avail

Spec

A-1



2. Inversions using lattice variables

Let X be a non-negative lattice variable with mean $u > 0$ and finite variance $\sigma^2 > 0$. Take X_1, X_2, \dots a sample from X . Define the sum

$$S_n(u) = X_1 + X_2 + \dots + X_n, \quad n \geq 1$$

and put

$$d_n(u, t) = P\{S_n(u) \leq [nt]\}.$$

By the weak law of large numbers we know that $S_n(u)/n \xrightarrow{P} u$ while $[nt]/n \rightarrow t$. Hence $d_n(u, t) \rightarrow 0$ if $t < u$, while $d_n(u, t) \rightarrow 1$ if $t > u$. For $u = t$ note that $\{[nu] - nu\}/\sqrt{n} \rightarrow 0$ while

$$d_n(u, u) = P\left\{\frac{S_n(u) - nu}{\sigma\sqrt{n}} \leq \frac{[nu] - nu}{\sigma\sqrt{n}}\right\} \rightarrow \frac{1}{2}$$

by the central limit theorem and the uniform convergence to the normal law [7, p. 139]. Hence

$$d_n(u, t) \rightarrow \begin{cases} 0 & t < u \\ \frac{1}{2} & t = u \\ 1 & t > u. \end{cases}$$

By Lebesgue's theorem and (2) for $0 < y_1 < y_2 < \infty$

$$(3) \quad \mu\{y_1; y_2\} = \lim_{n \rightarrow \infty} \int_0^\infty \{d_n(u, y_2) - d_n(u, y_1)\} d\mu(u)$$

is a potential inversion formula.

Moreover by a local limit theorem [4, p. 225] or [1, p. 233]

$$(4) \quad h_n(u, t) := \sigma \sqrt{2\pi n} P\{S_n(u) = [nt]\} \rightarrow \begin{cases} 0 & t \neq u \\ 1 & t = u \end{cases}.$$

By proper approximation (4) will suggest a function of the form

$$e_n(v) = ch_n(v, 1) \rightarrow \begin{cases} 0 & v \neq 1 \\ 1 & v = 1 \end{cases}$$

for a constant c ; this implies an inversion formula for the point mass at y , i.e.

$$(5) \quad \mu\{y\} = \lim_{n \rightarrow \infty} \int_0^{\infty} e_n\left(\frac{u}{y}\right) d\mu(y).$$

Before turning to examples let us introduce the abbreviation

$$(0 < y_1 < y_2 < \infty)$$

$$(6) \quad A_n = \{m: [ny_1] + 1 \leq m \leq [ny_2]\}.$$

Example 2.1. Poisson distribution and Laplace transform.

Put for the Laplace transform of μ ,

$$(7) \quad g(t) := \int_0^{\infty} e^{-tu} d\mu(u).$$

Then for $m \in \mathbb{N}$

$$(8) \quad (-1)^n g^{(n)}(t) = \int_0^{\infty} e^{-tu} u^n d\mu(u).$$

Let X be Poisson with mean u so that

$$d_n(u, t) = \sum_{m=0}^{[nt]} e^{-nu} \frac{(nu)^m}{m!}.$$

By (2) and (3) we get

$$\begin{aligned} \mu\{y_1; y_2\} &= \lim_{n \rightarrow \infty} \sum_{m \in A_n} \int_0^{\infty} e^{-nu} \frac{(nu)^m}{m!} d\mu(u) \\ &= \lim_{n \rightarrow \infty} \sum_{m \in A_n} \frac{(-nu)^m}{m!} g^{(m)}(n). \end{aligned}$$

This inversion formula can be found in [12, p. 295]. It can be fruitfully used in giving a probabilistic proof of Bernstein's theorem on completely monotone functions; see [2, p. 191].

For the point mass we get from (4)

$$h_n(u, t) = \sqrt{2\pi nu} e^{-nu} \frac{(nu)^{[nt]}}{[nt]!}.$$

Stirling's formula easily yields $c = 1$ and

$$e_n(v) = \exp - n\{v - 1 + \log v\}.$$

Collecting results we get

Theorem 2.1. Let μ be a bounded measure and $g(t)$ its Laplace transform.

Then for $0 < y_1 < y_2 < \infty$

$$\mu\{y_1; y_2\} = \lim_{n \rightarrow \infty} \sum_{m \in A_n} \frac{(-nu)^m}{m!} g^{(m)}(n);$$

and for $0 < y < \infty$

$$\mu\{y\} = \lim_{n \rightarrow \infty} \left(-\frac{e}{y}\right)^n g^{(n)}\left(\frac{n}{y}\right).$$

The latter formula occurs also in [12, p. 298].

Example 2.2. Pascal distribution and Stieltjes transform.

In the sequel we also deal with the (generalized) Stieltjes transform.

$$(9) \quad \phi_p(t) := \int_0^{\infty} (u+t)^{-p} d\mu(u)$$

where $p > 0$.

In evaluating successive derivatives of ϕ_p the next lemma will be useful.

Lemma 2.2. For $a, c \in \mathbb{N}$, $-p \notin \mathbb{N}_0$, $u > 0$, $t > 0$

$$(10) \quad D^a \{t^b D^c (u+t)^{-p}\} = \frac{\Gamma(p+a+c)}{\Gamma(p)} (-1)^{a+c} t^b (u+t)^{-p-a-c} F(-a, -b, -p-a-c+1; \frac{u+t}{t});$$

in particular

$$(11) \quad D^a \{t^{p+a+c-1} D^c (u+t)^{-p}\} = \frac{\Gamma(p+a+c)}{\Gamma(p)} (-1)^c t^{p+c-1} u^a (u+t)^{-p-a-c}.$$

Proof. Apply Leibniz's rule for differentiation of products; and then, for

$\alpha \in \mathbb{N}$

$$F(-\alpha, -\beta, -\gamma; x) = \sum_{m=0}^{\alpha} \frac{\binom{\alpha}{m} \binom{\beta}{m}}{\binom{\gamma}{m}} (-x)^m.$$

To prove (11), note that $F(-\alpha, \gamma, \gamma; x) = (1-x)^{\alpha}$ if $\alpha \in \mathbb{N}$ by [5, p. 1040]. \square

Let X be geometric with mean u . Then

$$d_n(u, t) = \sum_{m=0}^{[nt]} \binom{n+m-1}{m} p^m (1-p)^n$$

where $u = EX = \frac{1-p}{p}$ or $p = (u+1)^{-1}$. For the point mass we note that

$$h_n(u, t) = \sigma \sqrt{2\pi n} \binom{n+m-1}{m} \frac{u^m}{(1+u)^{n+m}}$$

where $m = [nt]$. Stirling's formula again applies to yield

$$(12) \quad e_n(v) = \left(\frac{2\sqrt{v}}{1+v}\right)^{2n}.$$

Theorem 2.3. Let μ be a bounded measure and $\phi_1(t)$ its (ordinary) Stieltjes transform. Then for $0 < y_1 < y_2 < \infty$

$$\begin{aligned} \mu\{y_1; y_2\} &= \lim_{n \rightarrow \infty} \frac{1}{\Gamma(n)} \sum_{m \in A_n} \frac{1}{m!} D^n \{t^{n+m-1} (-D)^{m-1} \phi_1(t)\}_{t=1} \\ &= \lim_{n \rightarrow \infty} \frac{1}{\Gamma(n)} \sum_{m \in A_n} \frac{(-1)^{m-1}}{m!} D^{n+m-1} \{t^n \phi_1(t)\}_{t=1}; \end{aligned}$$

for $0 < y < \infty$,

$$\mu\{y\} = \lim_{n \rightarrow \infty} \frac{2^{2n} y}{\Gamma(2n)} D^n \{y^{2n-1} (-D)^{n-1} \phi_1(y)\}.$$

Proof. From (3) and the current choice of d_n we find

$$\mu\{y_1; y_2\} = \lim_{n \rightarrow \infty} \sum_{m \in A_n} \binom{n+m-1}{m} \int_0^\infty \frac{u^m}{(u+1)^{m+n}} d\mu(u).$$

Put $a = n$, $b = n + m - 1$, $c = m - 1$, $p = 1$ in (11) to find the first

formula. To get the second, put $a = n + m - 1$, $b = n$, $c = 0$, $p = 1$ in (10) and recall that $F(a, b, c; x) = F(b, a, c; x)$.

For the point mass we get from (5) and (12)

$$\mu\{y\} = \lim_{n \rightarrow \infty} (4y)^n \int_0^{\infty} \frac{u^n}{(u+y)^{2n}} d\mu(u).$$

So we take $a = n$, $b = 2n - 1$, $c = n - 1$ and $p = 1$ in (11). □

Example 2.3. Binomial distribution and Stieltjes transform.

Let X be Bernoulli on 0 and 1 so that

$$P\{X = k\} = \begin{cases} 1 - v & k = 0 \\ v & k = 1. \end{cases}$$

Then

$$P\{S_n(v) \leq [nt]\} = \sum_{\ell=0}^{[nt]} \binom{n}{\ell} v^{\ell} (1-v)^{n-\ell}$$

suggests a potential inversion. However we should adapt it so that $v \in \mathbb{R}_+$ rather than $v \in [0, 1]$. So define

$$d_n(u, t) = \sum_{\ell=0}^{[n/(1+t)]} \binom{n}{\ell} u^{n-\ell} (1+u)^{-n}.$$

Theorem 2.4. Let μ be a bounded measure and $\phi_1(t)$ its (ordinary) Stieltjes transform. Then for $0 < y_1 < y_2 < \infty$

$$\mu\{y_1; y_2\} = \lim_{n \rightarrow \infty} \sum_{\ell \in B_m} \frac{1}{(n-1)!} \binom{n}{\ell} D^{n-\ell} \{t^{n-1} (-D)^{\ell-1} \phi_1(t)\}_{t=1}$$

$$= \lim_{n \rightarrow \infty} \sum_{\ell \in B_m} \frac{1}{(n-1)!} \binom{n}{\ell} D^{n-1} \{t^{n-\ell} \phi_1(t)\}_{t=1},$$

where $B_m = \{\ell: \lfloor \frac{n}{y_2+1} + 1 \rfloor \leq \ell \leq \lfloor \frac{n}{y_1+1} \rfloor\}$.

The proof is basically the same as that of the previous theorem.

3. Inversions using also densities

Let X_1, X_2, \dots be a sequence of independent non-negative random variables with common distribution F , mean 1 and finite variance. Let also Y_1, Y_2, \dots be such a sequence but with distribution G , and independent of the sequence $\{X_i\}$.

Define $S_n = X_1 + \dots + X_n$ and $T_n = Y_1 + \dots + Y_n$ and look at

$$d_n(u) = P\{S_n/T_n \leq u\};$$

by combining a weak law and the central limit theorem, one easily finds that

$$d_n(u) = \int_0^\infty F^{(n)}(ut) dG^{(n)}(t) \rightarrow \begin{cases} 0 & u < 1 \\ \frac{1}{2} & u = 1 \\ 1 & u > 1 \end{cases}$$

where $F^{(n)}(G^{(n)})$ is the n -fold convolution of $F(G)$ with itself. The particular case where $Y_i = 1$ is important and yields $d_n(u) = F^{(n)}(nu)$.

The potential inversion formula reads now

$$(13) \quad \mu\{y_1; y_2\} = \lim_{n \rightarrow \infty} \int_0^\infty \{d_n(\frac{u}{y_1}) - d_n(\frac{u}{y_2})\} d\mu(u)$$

while a point mass formula can be derived by using

$$e_n(v) = \frac{c}{\sqrt{n}} v d'_n(v)$$

if d_n is absolutely continuous.

Example 3.1. Exponential distribution and Laplace transform.

Here $Y_i = 1$ and X_i has an exponential density with mean 1. Hence

$$(14) \quad d_n(u) = \frac{1}{\Gamma(n)} \int_0^{nu} e^{-v} v^{n-1} dv$$

and so we have

Theorem 3.1. Let μ be a bounded measure and $g(t)$ its Laplace transform. Then for $0 < y_1 < y_2 < \infty$

$$(15) \quad \mu\{y_1; y_2\} = \lim_{n \rightarrow \infty} \int_{y_1}^{y_2} \frac{(-n)^n}{\Gamma(n)} \frac{g^{(n)}\left(\frac{n}{t}\right)}{t^{n+1}} dt.$$

Proof. From (13) and (14) we see that

$$\begin{aligned} \mu\{y_1; y_2\} &= \lim_{n \rightarrow \infty} \int_0^\infty d\mu(u) \int_{\frac{n}{y_2}}^{\frac{n}{y_1}} \frac{1}{\Gamma(n)} e^{-v} v^{n-1} dv \\ &= \lim_{n \rightarrow \infty} \int_{y_1}^{y_2} \frac{(-n)^n}{\Gamma(n)} \frac{dt}{t^{n+1}} \int_0^\infty e^{-u \frac{n}{t}} (-u)^n d\mu(u) \end{aligned}$$

by Fubini's theorem. □

The point mass formula coincides with that of th. 2.1.

The appealing form of (15) suggests that if the sequence

$$k_n(t) := \frac{(-n)^n}{\Gamma(n)} \frac{g^{(n)}\left(\frac{n}{t}\right)}{t^{n+1}}$$

converges a.e. on \mathbb{R}_+ and is bounded by an L_1 -function then μ is absolutely continuous and its derivative is given by the inversion

$$(16) \quad \frac{d\mu}{dt} = \lim_{n \rightarrow \infty} \frac{(-n)^n}{\Gamma(n)} \frac{g^{(n)}\left(\frac{n}{t}\right)}{t^{n+1}}.$$

The inversion formula (16) is well known [6,12]; its probabilistic proof due to Feller [3] has been the inspiring source for the current paper. Formula (16) has been used by Jagerman [9] to perform numerical inversion of the Laplace transform. For other applications see a paper by Vinogradov [13].

Example 3.2. Exponential distributions and Stieltjes transform.

Let $F(x) = G(x) = 1 - e^{-x}$; then $d_n(u)$ turns out to be a beta distribution on \mathbb{R}_+ , i.e.

$$d_n(u) = \frac{\Gamma(2n)}{\Gamma^2(n)} \int_0^u \frac{v^{n-1}}{(1+v)^{2n}} dv.$$

Theorem 3.2. Let μ be a bounded measure and $\phi_1(t)$ be its (ordinary) Stieltjes transform. Then for $0 < y_1 < y_2 < \infty$

$$\begin{aligned} \mu\{y_1; y_2\} &= \lim_{n \rightarrow \infty} \frac{1}{\Gamma^2(n)} \int_{y_1}^{y_2} D^n \{t^{2n-1} (-D)^{n-1} \phi_1(t)\} dt \\ &= \lim_{n \rightarrow \infty} \frac{(-1)^{n-1}}{\Gamma^2(n)} \int_{y_1}^{y_2} t^{n-1} D^{2n-1} \{t^n \phi_1(t)\} dt; \end{aligned}$$

for $0 < y < \infty$

$$\mu\{y\} = \lim_{n \rightarrow \infty} \frac{(-4y)^n}{\Gamma(2n)} D^{2n-1} \{y^n \phi_1(y)\}.$$

Proof. We easily get

$$\mu\{y_1; y_2\} = \lim_{n \rightarrow \infty} \frac{\Gamma(2n)}{\Gamma^2(n)} \int_{y_1}^{y_2} t^{n-1} dt \int_0^\infty \frac{u^n}{(u+t)^{2n}} d\mu(u).$$

We get the two formulas by applying lemma 2.2 with the two choices

$$a = n, b = 2n - 1, c = n - 1, p = 1 \text{ in (11)}$$

$$a = 2n - 1, b = n, c = 0, p = 1 \text{ in (10)}$$

as in the proof of th. 2.3.

For the point mass we get e_n as in (12); the resulting formula comes from the second choice above. \square

For the above formulas see [12, p. 350]. For applications, see [8].

The last example shows how an almost trivial probabilistic argument widens the applicability of the previous examples.

Example 3.3. Exponential distributions and generalized Stieltjes transforms.

Return to the previous example but look at S_{n+b} and T_{n+a+p} instead of S_n and T_n where $a, b \in \mathbb{Z}$ are fixed. Then

$$\lim_{n \rightarrow \infty} P\left\{\frac{S_{n+b}}{T_{n+a+p}} \leq u\right\} = \lim_{n \rightarrow \infty} P\left\{\frac{S_n}{T_n} \leq u\right\}.$$

Hence we have another candidate for d_n , i.e.

$$d_n(u) = \frac{\Gamma(2n+a+b+p)}{\Gamma(n+a+p)\Gamma(n+b)} \int_0^u \frac{v^{n+b-1}}{(1+v)^{2n+a+b+p}} dv.$$

By the usual procedure we obtain also

$$e_n(v) = v^{\frac{b-a-p}{2}} \left\{\frac{2\sqrt{v}}{1+v}\right\}^{2n+a+b+p}.$$

If we now go through the same routine as in example 3.2 with a replaced by $n + b$ and c by $n + a$ we obtain the following results of E.R. Love and A. Byrne [10,11]. We note that we have to restrict p to \mathbb{N}_0 in view of the

probabilistic set up.

Theorem 3.3. Let μ be a bounded measure ; let $\phi_p(t)$ be its generalized Stieltjes transform. Let $a, b \in \mathbb{Z}$. Then for $0 < y_1 < y_2 < \infty$

$$\mu\{y_1; y_2\} = \lim_{n \rightarrow \infty} \frac{\Gamma(p)}{\Gamma(n+a+p)\Gamma(n+b)} \int_{y_1}^{y_2} D^{n+b} \{ t^{2n+a+b+p-1} (-D)^{n+a} \phi_p(t) \} dt;$$

for $0 < y < \infty$

$$\mu\{y\} = \lim_{n \rightarrow \infty} \frac{2^{2n+a+b+p} \Gamma(p)}{\Gamma(2n+a+b+p)} y D^{n+b} \{ y^{2n+a+b+p-1} (-D)^{n+a} \phi_p(y) \}.$$

Note that by using different random variables one can get a variety of real inversion formulas for the same Stieltjes transform ϕ_1 . Needless to point out how th. 3.2 follows from the last result.

4. Remarks.

1. The main drawback of the method used in this paper is that the relevant integral transform has to be recovered from a potential inversion formula.

2. The main virtue of the paper lies in singling out the relevance of the asymptotic behavior of the sequence d_n for $n \rightarrow \infty$; probabilistic arguments often provide this behavior immediately.

3. There exist additive counterparts for the formula (13). For if

$$d_n(v) \rightarrow \begin{cases} 0 & v < 0 \\ \frac{1}{2} & v = 0 \\ 1 & v > 0 \end{cases}$$

where $v \in \mathbb{R}$, then a potential inversion formula for a transform on \mathbb{R} is

$$\mu\{y_1; y_2\} = \lim_{n \rightarrow \infty} \int_{\mathbb{R}} \{d_n(u - y_1) - d_n(u - y_2)\} d\mu(u).$$

The best known example is $d_n(v) = \frac{1}{\pi} \int_{-\infty}^{nv} \frac{\sin u}{u} du$ leading to the inversion formula for the Fourier transform

$$\mu\{y_1; y_2\} = \lim_{n \rightarrow \infty} \int_{y_1}^{y_2} dx \left\{ \frac{1}{2\pi} \int_{-n}^n e^{-izx} \phi(z) dz \right\}$$

where $\phi(z) = \int_{\mathbb{R}} e^{izu} d\mu(u)$.

Another example is $d_n(v) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\sqrt{nv}} e^{-\frac{1}{2}x^2} dx$ leading to a real inversion formula for the Gauss-Weierstrasz transform [6]

$$w_\lambda(t) = \sqrt{\frac{t}{2\pi}} \int_{-\infty}^{\infty} \exp - \frac{\lambda}{2}(u - t)^2 d\mu(u).$$

References

- [1] BREIMAN, L.: Probability, Addison-Wesley, Reading Mass., 1968.
- [2] CHUNG, K.L.: A Course in Probability Theory, 2d. ed. Academic Press, New York, 1974.
- [3] FELLER, W.: An Introduction to Probability Theory and Its Applications, Vol. II, Wiley, New York, 1971.
- [4] GNEDENKO, B.V. & KOLMOGOROV, A.N.: Limit Distributions for Sums of Independent Random Variables, Addison-Wesley, Reading, Mass., 1968.
- [5] GRADHSTEIN, I.S. & RYZHIK, I.M.: Tables of Integrals, Series, and Products, Academic Press, New York, 1975.
- [6] HIRSCHMAN, I.I. & WIDDER, D.V.: The Convolution Transform, Princeton Univ. Press, Princeton, New Jersey, 1955.
- [7] IBRAGIMOV, I.A. & LINNIK, Y.V.: Independent and Stationary Sequences of Random Variables, Noordhoff, Groningen, 1971.
- [8] ISMAIL, M.E.H.: Bessel functions and the infinite divisibility of the Student t-distribution, Ann. Probability, 1977, 5, 582-585.
- [9] JAGERMAN, D.L.: An inversion technique for the Laplace transform with application to approximation, Bell System Tech. J., 1978, 57, 669-710.
- [10] LOVE, E.R. & BYRNE, A.: Real inversion theorems for generalized Stieltjes transforms, J. London Math. Soc., (2), 1980, 22, 285-306.
- [11] LOVE, E.R. & BYRNE, A.: Real inversion theorems for generalized Stieltjes transforms II, Math. Proc. Camb. Phil. Soc., 1982, 92, 275-291.
- [12] WIDDER, D.V.: The Laplace Transform, Princeton Univ. Press, Princeton, New Jersey, 1946.
- [13] VINOGRADOV, O.P.: On applications of an inversion formula for the Laplace transform, Th. Probability Appl., 1979, 21, 836-839.

END

FILMED

1-86

DTIC